




MoreNova Ortho Report 05/05/2024

Nine patients have undergone treatment until today. The patients did not take any NSAIDs (nonsteroidal anti-inflammatory drugs) and did not receive any other kind of treatment. Each patient has received one session per week, with 1000-1500 shocks applied per session. Below is a table showing the pain scale scores before and after the Li-shockwaves treatment.

Patient number	Diagnosis	Age	Sex	Pain Pre	Pain Post	Number of sessions
1	Epitrocleitis	32	M	8	3	5
2	Epitrocleitis	55	F	9	1	3
3	Achilles tendonitis	59	F	7	5	6
4	Patellar tendonitis	41	M	8	3	3
5	Biceps tendinitis	51	M	8	5	4
6	Epicondylitis	44	M	9	4	5
7	Trochanteritis	68	F	7	3	5
8	Patellar tendonitis	59	M	8	4	6
9	Plantar fasciitis	71	F	9	7	6



Dr. Osvaldo Patiño

Director of Specialization in Kinesiology and Sports
Physiotherapy at Universidad del Gran Rosario

<https://www.researchgate.net/profile/Osvaldo-Patino>

<https://www.linkedin.com/in/osvaldo-pati%C3%B1o-408691179?originalSubdomain=ar>

<https://www.instagram.com/dr.osvaldo.patino>